

[54] OPTHALMOLOGICAL PROCEDURES

[75] Inventor: Lawrence G. Pape, Scarsdale, N.Y.

[73] Assignee: Ophthalmic Systems, Inc., Long Island City, N.Y.

[21] Appl. No.: 198,290

[22] Filed: Oct. 20, 1980

[51] Int. Cl.³ A61B 17/00; A61K 31/70

[52] U.S. Cl. 128/276; 424/180

[58] Field of Search 128/276

[56] References Cited

U.S. PATENT DOCUMENTS

4,141,973 2/1979 Balazs 424/180

OTHER PUBLICATIONS

Healon, Product Information, Pharmacia.

Pape, L. G., The Use of Sodium Hyaluronate, *Ophthalmology*, vol. 87, 1980.

Binkhorst, C. D., American Intra-Ocular Implant Society, *Journal*, vol. 6 (4), Oct. 1980, pp. 340-341.

Primary Examiner—Robert W. Michell

Assistant Examiner—Nancy A. B. Swisher

Attorney, Agent, or Firm—Joseph Patrick Burke

[57] ABSTRACT

This disclosure is directed to improved ophthalmological surgical procedures on the human eyeball characterized by initially utilizing ultra-pure hyaluronic acid in the form of a solution containing the sodium salt thereof, in a sufficient concentration thereof to protect delicate eye tissue and cells, e.g., 0.5 to 1.0 wt. % and higher, and thereafter, significantly diluting the concentration thereof in the aqueous media (natural and/or synthetic) within the anterior segment of the eye viz, in situ, so as to arrive at a concentration of said sodium salt of hyaluronic acid of less than 0.5 wt. % and more usually from about 0.1 to about 0.3 wt. % prior to closure. Hence a significant dilution from the initial higher concentrations by a factor of at least about 2:1 and more usually from about 3:1 to about 10:1 is effected. Hence the concentration of said sodium hyaluronate in the natural and/or synthetic aqueous media left within the eyeball at the surgical site after closure of the procedure ranges from about 0.1 to about 0.3 wt. % in the anterior chamber.

21 Claims, 2 Drawing Figures

